










The doses include in this chart are clinician recommended. **DO NOT give more than 5 doses in 24 hour or every 4 to 6 hours**

*dose in some dosage forms does not provide the same exact dose, it will be rounded to the nearest possible dosage listed in ()

Active ingredient: Acetaminophen	6-9 lbs (3-4 kg)	10-12 lb (4.5 -5.5 kg)	13-17 lbs (6-7.5 kg)	18-23 lbs (8-10 kg)	24-35 lbs (11-15 kg)	36-46 lbs (16-21 kg)	47-58 lbs (22-26 kg)	59-70 lbs (27-31 kg)	71-93 lbs (32-42 kg)	94 Lbs + (43 kg+)
Medication Dosage Forms:										
Dose	40 mg	64 mg	80 mg	120 mg	160 mg	240 mg	320 mg*	400 mg*	480 mg*	640 mg*
Oral Suspension 160 mg/ 5mL <ul style="list-style-type: none"> Infant Acetaminophen Children Acetaminophen 	1.25 mL 	2 mL ♦ 	2.5 mL 	3.75 mL 	5 mL  or 	7.5 mL 	10 mL 	12.5 mL 	15 mL 	20 mL  
Chewable Tablet 160 mg each					1 chewable tablet 	1 ½ chewable tablets 	2 chewable tablets 	2 ½ chewable tablets 	3 chewable tablets 	4 chewable tablets 
Regular Strength Tablet 325 mg each DO NOT USE: Extra Strength 500 mg Tablet <i>or</i> Extended-release 650mg tablet							1 tablet  (325 mg)	1 tablet  (325 mg)	1½ tablets  (487.5 mg)	2 tablets  (650 mg)
Suppository 80 mg each <ul style="list-style-type: none"> Infant Fever All 			1 suppository 	1 ½ suppositories 	2 suppositories 	3 suppositories 				
Suppository 120 mg each <ul style="list-style-type: none"> Children Fever All 				1 suppository 		2 suppositories 		3 suppositories  (360 mg)		
Suppository 325 mg each <ul style="list-style-type: none"> Jr. Fever All 							1 suppository  (325 mg)	1 suppository  (325 mg)	1 suppository  (325 mg)	2 suppositories  (650 mg)

♦ Measuring syringe may not come with acetaminophen packaging, ask pharmacy for 3 mL oral syringe for measuring